

Light Pseudoscalar Mesons Decays at JLab

Liping Gan

University of North Carolina Wilmington

Outline

- Introduction of Physics
- PrimEx Primakoff Program
- JLab Eta Factory (JEF) at GlueX
- Summary

Thanks for support by NSF PHY-1812396 and PHY-2111181.

Open Questions in Modern Physics

- What is the origin of QCD confinement?
- How did the visible mass emerge in the early universe?
- What is the cause of the matter-antimatter asymmetry in the universe?
- What is the nature of dark matter?

Light pseudoscalar mesons offer a sensitive tool to explore these fundamental questions.

Low-Energy QCD Symmetries and Light Mesons

- QCD Lagrangian in Chiral limit ($m_q \rightarrow 0$) is invariant under:

$$SU_L(3) \times SU_R(3) \times U_A(1) \times U_B(1)$$

- Chiral symmetry $SU_L(3) \times SU_R(3)$ spontaneously breaks to $SU(3)$

- 8 Goldstone Bosons (GB)


- $U_A(1)$ is explicitly broken:

(Chiral anomalies)

- Non-zero mass of η_0
 - $\Gamma(\pi^0 \rightarrow \gamma\gamma), \Gamma(\eta \rightarrow \gamma\gamma), \Gamma(\eta' \rightarrow \gamma\gamma)$

- $SU_L(3) \times SU_R(3)$ and $SU(3)$ are explicitly broken:

- GB are massive
 - Mixing of π^0, η, η'




The π^0, η, η' system provides a rich laboratory to study the symmetry structure of low-energy QCD.

What is the origin of visible mass?

Mass-generating mechanisms:

- Higgs boson, alone is responsible for $<2\%$ of the visible mass in the universe.
- Emergent Hadron Mass (EHM) and its constructive interference with Higgs-boson account for $>98\%$ of the visible mass.



Few Body Syst. 63 (2022) 2,42
Few Body Syst. 65 (2024) 2,60

Complementary to proton, pseudoscalar mesons offer a unique opportunity to study the interference between two known mass generating mechanisms.

Discrete Symmetries

Class	Violated	Conserved	Interaction
0		C, P, T, CP, CT, PT, CPT	strong, electromagnetic
I	C, P, CT, PT	T, CP, CPT	(weak, with no KM phase or flavor-mixing)
II	P, T, CP, CT	C, PT, CPT	
III	C, T, PT, CP	P, CT, CPT	
IV	C, P, T, CP, CT, PT	CPT	weak

Class II: P-, CP-violation

- QCD θ -term
- Examples: $\eta^{(\prime)} \rightarrow 2\pi$, $\eta^{(\prime)} \rightarrow \pi^+\pi^-\gamma^{(*)}$
- Strong constraints from EDM measurements

Class III: C-, CP-violation


- A C- and T-violating, and P-conserving interaction was proposed by Bernstein, Feinberg and Lee, but little theoretic progress for a long time. [Phys. Rev.,139, B1650 \(1965\)](#)
- Recent new theoretic development (presented by B. Kubis this morning).
- Electroweak radiative corrections mix class II and III, but much weaker EDM constraints.
- Examples: $\eta^{(\prime)} \rightarrow 3\gamma$, $\eta^{(\prime)} \rightarrow \pi^0\gamma^{(*)}$...

Class III has much weaker experimental constraint, offer an opportunity for experimental search of new physics in η decays.

Class III: C- and CP-Violation in $\eta^{(\prime)} \rightarrow \pi^+\pi^-\pi^0$, $\eta' \rightarrow \pi^+\pi^-\eta$



- Dalitz plot decomposition (central fit result)

$$|\mathcal{M}_c|^2 \approx |\mathcal{M}_1^C|^2 + 2\text{Re} [\mathcal{M}_1^C (\mathcal{M}_0^\phi)^*] + 2\text{Re} [\mathcal{M}_1^C (\mathcal{M}_2^\phi)^*]$$



- \mathcal{M}_0^ϕ and \mathcal{M}_2^ϕ lead to different interference patterns
- CP-violation from these processes is not bounded by EDM.
- Complementary to nEDM searches even in the case of T and P odd observables, since the flavor structure of the η is different from the nucleon.

BSM Physics in Dark Sector



Dark Sector

- New gauge forces, bosons and fermions beyond SM.
- The stability of dark matter can be explained by the dark charge conservation.

Portals Coupling SM and Dark Sector

Standard Model:
 $SU(3) \times SU(2) \times U(1)$



Dark Sector:
Gauge Interactions?
Dark matter?

vector:

- Leptophobic vector B'

$$\eta, \eta' \rightarrow B' \gamma \rightarrow \pi^0 \gamma \gamma, \quad (0.14 < m_{B'} < 0.62 \text{ GeV});$$

$$\eta' \rightarrow B' \gamma \rightarrow \pi^+ \pi^- \pi^0 \gamma, \quad (0.62 < m_{B'} < 1 \text{ GeV}).$$

- X boson or dark photon: $\eta, \eta' \rightarrow X \gamma \rightarrow e^+ e^- \gamma$

scalar S: $\eta \rightarrow \pi^0 S \rightarrow \pi^0 \gamma \gamma, \pi^0 e^+ e^-, \quad (10 \text{ MeV} < m_S < 2m_\pi);$

$$\eta, \eta' \rightarrow \pi^0 S \rightarrow 3\pi, \eta' \rightarrow \eta S \rightarrow \eta \pi \pi, \quad (m_S > 2m_\pi).$$

Fermion: $\eta \rightarrow \pi^0 H,$

$$\text{with } H \rightarrow \nu N_2, N_2 \rightarrow h' N_1, h' \rightarrow e^+ e^-$$

Portals:

vector $\kappa B^{\mu\nu} V_{\mu\nu}$


Scalar $H^+ H (\epsilon S + \lambda S^2)$

Fermion $\xi L H N$

ALP $c_{\gamma\gamma} \frac{\alpha}{4\pi} \frac{a}{f} F_{\mu\nu} \tilde{F}^{\mu\nu} + c_{GG} \frac{\alpha_s}{4\pi} \frac{a}{f} G_{\mu\nu}^a \tilde{G}^{a,\mu\nu}$

Axion-Like Particles (ALP): $\eta, \eta' \rightarrow \pi \pi a \rightarrow \pi \pi \gamma \gamma, \pi \pi e^+ e^-$


Landscape of BSM Physics Search




Rept. Prog. Phys. 79, no.12, 124201

Complementary to other types of experiments, pseudoscalar mesons offer unique sensitivity for sub-GeV new physics that are flavor-conserving and light quark-coupling.

Jefferson Lab



Primakoff Effect




$$\frac{d\sigma_{\text{Pr}}}{d\Omega} = \boxed{\Gamma_{\gamma\gamma}} \frac{8\alpha Z^2}{m_\pi^3} \frac{\beta^3 E^4}{Q^4} |F_{e.m.}(Q)|^2 \sin^2 \theta_\pi$$

- Peaked at very small forward angle: $\langle \theta_{\text{Pr}} \rangle_{\text{peak}} \propto \frac{m^2}{2E^2}$
- Beam energy sensitive:

$$\left\langle \frac{d\sigma_{\text{Pr}}}{d\Omega} \right\rangle_{\text{peak}} \propto \frac{E^4}{m^3}, \quad \int d\sigma_{\text{Pr}} \propto \frac{Z^2}{m^3} \log E$$

$$\boxed{\langle \theta_{\text{Pr}} \rangle_{\text{peak}} \propto \frac{m^2}{2E^2} \quad \langle \theta_{\text{NC}} \rangle_{\text{peak}} \propto \frac{2}{E \cdot A^{1/3}}}$$

- Coherent process



- The higher beam energy is, the higher Primakoff cross section and the better separation of Primakoff from the nuclear backgrounds.
- A higher beam energy is more important for more massive particle

PrimEx Primakoff Program at JLab 6 & 12 GeV


Precision measurements of electromagnetic properties of π^0 , η , η' via Primakoff effect

a) Two-Photon Decay Widths:

- 1) $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ @ 6 GeV
- 2) $\Gamma(\eta \rightarrow \gamma\gamma)$
- 3) $\Gamma(\eta' \rightarrow \gamma\gamma)$

Input to Physics:

- precision tests of chiral symmetry and anomalies
- light quark mass ratio
- η - η' mixing angle
- input to calculate HLbL in $(g-2)_\mu$
- origin of the visible mass



b) Transition Form Factors

at Q^2 of 0.001-0.3 GeV^2/c^2 :

$$F(\gamma\gamma^* \rightarrow \pi^0), F(\gamma\gamma^* \rightarrow \eta), F(\gamma\gamma^* \rightarrow \eta')$$

Input to Physics:

- π^0, η and η' electromagnetic interaction radii
- is the η' an approximate Goldstone boson?
- input to calculate HLbL in $(g-2)_\mu$
- origin of the visible mass

Status of Primakoff Program at JLab 6 & 12 GeV

Precision measurements of electromagnetic properties of π^0 , η , η' via Primakoff effect

a) Two-Photon Decay Widths:


1) $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ @ 6 GeV (in Hall B)

2) $\Gamma(\eta \rightarrow \gamma\gamma)$

3) $\Gamma(\eta' \rightarrow \gamma\gamma)$

Input to Physics:


- precision tests of chiral symmetry and anomalies
- determination of light quark mass ratio
- η - η' mixing angle
- input to calculate HLbL in $(g-2)_\mu$



- The chiral anomaly prediction **is exact** for massless quarks:


$$\Gamma(\pi^0 \rightarrow \gamma\gamma) = \frac{m_{\pi^0}^3 \alpha^2 N_c^2}{576 \pi^3 F_{\pi^0}^2} = 7.750 \pm 0.016 \text{ eV}$$

- $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ is one of the few quantities in confinement region that QCD can calculate precisely at ~1% level to higher orders!



Status of Primakoff Program at JLab 6 & 12 GeV (cont.)

Precision measurements of electromagnetic properties of π^0 , η , η' via Primakoff effect



a) Two-Photon Decay Widths:

- 1) $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ @ 6 GeV
- 2) $\Gamma(\eta \rightarrow \gamma\gamma)$
- 3) $\Gamma(\eta' \rightarrow \gamma\gamma)$

Input to Physics:

- precision tests of chiral symmetry and anomalies
- determination of light quark mass ratio
- η - η' mixing angle
- input to calculate HLbL in $(g-2)_\mu$


$$\frac{d\sigma_{Pr}}{d\Omega} = \Gamma_{\gamma\gamma} \frac{8\alpha Z^2 \beta^3 E^4}{m_\eta^3 Q^4} |F_{e.m.}(Q^2)|^2 \sin^2 \theta_\eta$$

On-Going PrimEx-eta experiment (in Hall D)

- A full data set was completed via three run periods in 2019, 2021 and 2022.
- Data analysis is in progress.
(see talk by A. Somov on Tue)

Physics for $\Gamma(\eta \rightarrow \gamma\gamma)$ Measurement

Resolve long standing discrepancy between previous collider and Primakoff measurements:



- **Extract η - η' mixing angle**
- **Improve calculation of the η -pole contribution to Hadronic Light-by-Light (HLbL) scattering in $(g-2)_\mu$**
- **Improve all partial decay widths in the η -sector**


Precision Determination Light Quark Mass Ratio

A clean probe for quark mass ratio: $Q^2 = \frac{m_s^2 - \hat{m}^2}{m_d^2 - m_u^2}$, where $\hat{m} = \frac{1}{2}(m_u + m_d)$


➤ $\eta \rightarrow 3\pi$ decays through isospin violation: $A = (m_u - m_d)A_1 + \alpha_{em}A_2$

➤ α_{em} is small

➤ Amplitude: $A(\eta \rightarrow 3\pi) = \frac{1}{Q^2} \frac{m_K^2}{m_\pi^2} (m_\pi^2 - m_K^2) \frac{M(s, t, u)}{3\sqrt{3}F_\pi^2}$




Space-Like Transition Form Factors ($Q^2 : 0.001\text{-}0.3 \text{ GeV}^2/\text{c}^2$)





- Direct measurement of slopes


- Interaction radii:
 $F_{\gamma\gamma^*P}(Q^2) \approx 1 - \frac{1}{6} \cdot \langle r^2 \rangle_P Q^2$
- ChPT for large N_c predicts relation between the three slopes. Extraction of $O(p^6)$ low-energy constant in the chiral Lagrangian



- Input for hadronic light-by-light calculations in muon ($g-2$)



Phys.Rev.D65,073034




New opportunities with JLab 22 GeV Upgrade

1. The first π^0 Primakoff production off an electron target to measure $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ and $F(\gamma\gamma^* \rightarrow \pi^0)$.
2. Improve the precisions of η/η' Primakoff production off nuclear targets.
3. Search for new sub-GeV gauge bosons (scalars and pseudoscalars) via the Primakoff production:
 - Strong CP and Hierarchy problems
 - $(g - 2)_\mu$ and puzzle of proton charge radius
 - Portals coupling SM to the dark sector:

$$H^+ H (\epsilon S + \lambda S^2) \quad c_{\gamma\gamma} \frac{\alpha}{4\pi} \frac{a}{f} F_{\mu\nu} \tilde{F}^{\mu\nu} + c_{GG} \frac{\alpha_s}{4\pi} \frac{a}{f} G_{\mu\nu}^a \tilde{G}^{a,\mu\nu}$$


Advantages of the π^0 Primakoff Production off an Electron

PrimEx-II: $\gamma + {}^{28}\text{Si} \rightarrow \pi^0 + {}^{28}\text{Si}$



$$\frac{d\sigma_{\text{Pr}}}{d\Omega} = \boxed{\Gamma_{\gamma\gamma}} \frac{8\alpha Z^2}{m_\pi^3} \frac{\beta^3 E^4}{Q^4} |F_{e.m.}(Q)|^2 \sin^2 \theta_\pi$$

$E_\gamma: 4.45\text{--}5.30 \text{ GeV}$




Main challenges for the nuclear target:

- Nuclear backgrounds
- Nuclear effects
- No recoil detection

Advantages of an electron target:


- Eliminate all nuclear backgrounds
- A point-like electron target to eliminate nuclear effects
- Recoiled electron detection



$$\frac{d\sigma_{\text{Pr}}}{d\Omega} = \Gamma_{\gamma\gamma} \frac{8\alpha}{m_\pi^3} \frac{\beta^3 E^4}{Q^4} \sin^2 \theta_\pi$$

Measurement	Reaction	E_{th} (GeV)
$\Gamma(\pi^0 \rightarrow \gamma\gamma)$	$\gamma + e \rightarrow \pi^0 + e$	18.0
$F(\gamma^*\gamma \rightarrow \pi^0)$	$e + e \rightarrow \pi^0 + e + e$	18.1


Projected $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ at JLab 22 GeV with an Electron Target





Theory and Experiments

Improve Primakoff Measurements of η/η' with nuclear targets


$E_\gamma = 10$ GeV



$E_\gamma = 20$ GeV



Search for sub-GeV Scalar and Pseudoscalar via Primakoff Effect



$$\mathcal{L}_{\text{eff}} \supset \frac{c_\gamma}{4\Lambda} a F^{\mu\nu} \tilde{F}_{\mu\nu}$$

$$\frac{d\sigma_{Pr}}{d\Omega} \sim \frac{c_\gamma^2 \alpha Z^2}{8\pi\Lambda^2} \cdot \frac{\beta^3 E^4}{Q^4} \cdot |F_{e.m.}(Q)|^2 \sin^2 \theta_a$$



The Primakoff signal dominates in the forward angles



Minimizing the QCD backgrounds

Favorable experimental condition:


- A high energy beam
- A high Z nuclear target



PrimEx I

Phys.Rev.Lett. 106 (2011) 162303


Projected Reach for a ALP at JLab 22 GeV



$$\gamma + Pb \rightarrow a + Pb$$

$$a \rightarrow \gamma\gamma$$

JLab Eta Factory (JEF) Experiment at GlueX



- ◆ Simultaneously produce η/η' on LH_2 target with **8.4-11.7 GeV tagged photon beam** via $\gamma + p \rightarrow \eta/\eta' + p$
- ◆ Reduce non-coplanar backgrounds by **detecting recoil protons** with GlueX detector
- ◆ Upgraded Forward Calorimeter with **High resolution, high granularity PbWO₄ insertion (FCAL-II)** to detect multi-photons from the η/η' decays
- ◆ The GlueX detector will detect the charged products from the η/η' decays

Uniqueness of JEF Experiment

1. Suppressed backgrounds in rare neutral decays comparing to the other experiments using:
 - a) η/η' energy boost;
 - b) upgraded FCAL-II;
 - c) recoil detection
2. Capability of running in parallel with GlueX and other experiments in Hall D
→ potential for a high-statistics data set
3. Simultaneously produce tagged η and η' with similar rates ($\sim 5 \times 10^5$ per day)

Main JEF Physics Objectives

1. Search for sub-GeV hidden bosons

vector:

- Leptophobic vector B'

$$\eta, \eta' \rightarrow B' \gamma \rightarrow \pi^0 \gamma \gamma, (0.14 < m_{B'} < 0.62 \text{ GeV});$$

$$\eta' \rightarrow B' \gamma \rightarrow \pi^+ \pi^- \pi^0 \gamma, (0.62 < m_{B'} < 1 \text{ GeV}).$$

- Hidden or dark photon: $\eta, \eta' \rightarrow X \gamma \rightarrow e^+ e^- \gamma$.

scalar S: $\eta \rightarrow \pi^0 S \rightarrow \pi^0 \gamma \gamma, \pi^0 e^+ e^-, (10 \text{ MeV} < m_S < 2m_\pi);$

$$\eta, \eta' \rightarrow \pi^0 S \rightarrow 3\pi, \eta' \rightarrow \eta S \rightarrow \eta \pi \pi, (m_S > 2m_\pi).$$

Axion-Like Particles (ALP): $\eta, \eta' \rightarrow \pi \pi a \rightarrow \pi \pi \gamma \gamma, \pi \pi e^+ e^-$

2. Directly constrain CVPC new physics: $\eta^{(\prime)} \rightarrow 3\gamma, \eta^{(\prime)} \rightarrow 2\pi^0 \gamma, \eta^{(\prime)} \rightarrow \pi^+ \pi^- \pi^0$


3. Precision tests of low-energy QCD:

- Interplay of VMD & scalar dynamics in ChPT: $\eta \rightarrow \pi^0 \gamma \gamma, \eta' \rightarrow \pi^0 \gamma \gamma$
- Transition Form Factors of $\eta^{(\prime)}$: $\eta^{(\prime)} \rightarrow e^+ e^- \gamma$

4. Improve the light quark mass ratio via Dalitz distributions of $\eta \rightarrow 3\pi$

Example of a Key Channel: $\eta \rightarrow \pi^0 \gamma \gamma$

1. New physics:



Portal: ($n = 4$)


vector	$\kappa B^{\mu\nu} V_{\mu\nu}$
Scalar	$H^+ H (\varepsilon S + \lambda S^2)$
fermion	ξLHN

❖ Search for sub-GeV gauge bosons

- A leptophobic **vector** B' :
 $\eta \rightarrow \gamma B', B' \rightarrow \pi^0 \gamma$ PR, D89, 114008
- An electrophobic **scalar** Φ' :
 $\eta \rightarrow \pi^0 \Phi', \Phi' \rightarrow \gamma \gamma$

PRL 117, 101801 (2016); PL B740, 61 (2015)

2. Confinement QCD:




Gasser, Leutwyler 84; Ecker, Gasser, Pich, de Rafael 1989; Donoghue, Ramirez, Valencia 1989

❖ A rare window to probe interplay of VMD & scalar resonance in ChPT


JEF Experimental Reach for B'

A search for a leptophobic dark B' boson coupled to baryon number is complementary to ongoing searches for a dark photon

$$\eta \rightarrow B' \gamma \rightarrow \pi^0 \gamma \gamma$$




PL, B221, 80 (1989)
PR,D89,114008




Projected JEF on SM Allowed $\eta \rightarrow \pi^0 \gamma \gamma$

J.N. Ng and D.J. Peters, Phys. Rev. D47, 4939



χ PTh by Oset et al., Phys. Rev. D77, 073001



We measure both BR and Dalitz distribution

- ◆ model-independent determination of two LEC's of the $O(p^6)$ counter-terms
- ◆ probe the role of scalar resonances to calculate other unknown $O(p^6)$ LEC's

J. Bijnens, talk at AFCL workshop

Test Charge Conjugation Invariance

- ◆ C is maximally violated in the weak force and is well tested.
- ◆ Assumed in SM for electromagnetic and strong forces, but **it is not experimentally well tested**
(current direct constraint: $\Lambda \geq 1 \text{ GeV}$)

C Violating η neutral decays

Mode	Branching Ratio (upper limit)	No. γ 's
3γ	$< 1.6 \cdot 10^{-5}$	3
$\pi^0\gamma$	$< 9 \cdot 10^{-5}$	
$2\pi^0\gamma$	$< 5 \cdot 10^{-4}$	5
$3\gamma\pi^0$	Nothing published	
$3\pi^0\gamma$	$< 6 \cdot 10^{-5}$	7
$3\gamma 2\pi^0$	Nothing published	


Experimental Improvement on C-violating $\eta \rightarrow 3\gamma$

- ◆ SM contribution:
 $\text{BR}(\eta \rightarrow 3\gamma) < 10^{-19}$ via P-violating weak interaction.

- ◆ A calculation due to new physics by Tarasov suggests:

$$\text{BR}(\eta \rightarrow 3\gamma) < 10^{-2}$$

Sov.J.Nucl.Phys.,5,445 (1967)




Improve BR upper limit by one order of magnitude to directly tighten the constraint on CVPC new physics

Status of the JEF Experiment

1. Developed an upgraded FCAL-II with a PbWO_4 insert.

- 1596 PbWO_4 modules are developed to replace ~ 400 Pb-glass modules.
- Installation of the upgraded FCAL-II has been on-going since Mar 2023 and will be completed by the end of 2024.
- Over 40 undergraduate students from 11 institutes were trained by involving in this project.




Undergraduate workforce



May 2023



Oct 6, 2023



2. Commissioning of FCAL-II and data taking with FCAL-II are scheduled to start in Jan 2025.

Summary

- ◆ Light pseudoscalar mesons offer a sensitive probe to test fundamental symmetries and to search for new physics beyond the standard model.
- ◆ PrimEx Primakoff program
 - has been in progress @ 6&12 GeV
 - ✓ The published PrimEx result on the π^0 lifetime provides a stringent test of low-energy QCD.
 - ✓ Data collection on $\Gamma(\eta \rightarrow \gamma\gamma)$ was completed in 2022 and data analysis is in progress.
 - ✓ A new experiment on $F(\pi^0 \rightarrow \gamma^* \gamma)$ off a nuclear target is on the way.
- future JLab 22 GeV upgrade will offer new opportunities
 - ✓ New generation of Primakoff experiments on $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ and $F(\gamma^* \gamma \rightarrow \pi^0)$ off an atomic electron target.
 - ✓ Improve measurements of more massive particles, such as η and η' , off nuclear targets.
 - ✓ Search for new sub-GeV gauge bosons (scalars and pseudoscalars).
- ◆ The JEF experiment will start data collection in Jan 2025 using newly upgraded FCAL-II calorimeter with a PbWO4 insert.
 - ✓ Search for sub-GeV hidden bosons: vector, scalar, and ALP
 - ✓ Directly constrain CVPC new physics
 - ✓ Precision tests of low-energy QCD: the role of scalar dynamics in ChPT; transition form factors of η/η' to calculate HLbL contributions in $(g-2)_\mu$